

EQUIVALENCE OF MIRROR FAMILIES CONSTRUCTED FROM TORIC DEGENERATIONS OF FLAG VARIETIES

JOE RUSINKO

ABSTRACT. Batyrev (et. al.) constructed a family of Calabi-Yau varieties using small toric degenerations of the full flag variety G/B . They conjecture this family to be mirror to generic anti-canonical hypersurfaces in G/B . Recently Alexeev and Brion, as a part of their work on toric degenerations of spherical varieties, have constructed many degenerations of G/B . For any such degeneration we construct a family of varieties, which we prove coincides with Batyrev's in the small case. We prove that any two such families are birational, thus proving that mirror families are independent of the choice of degeneration. The birational maps involved are closely related to Berenstein and Zelevinsky's geometric lifting of tropical maps to maps between totally positive varieties.

1. INTRODUCTION

Generic elements of the anti-canonical class of a flag variety G/P are smooth Calabi-Yau varieties. We restrict our attention to G of type A_n (i.e. $G = SL_{n+1}$), and work over the ground field $\mathbb{C}$. For any parabolic subgroup P , Batyrev, Ciocan-Fontanine, Kim, and van Straten [BCFKvS00] use Gonciulea and Lakshmibai's [GL96] toric degeneration of G/P , to construct a family of CYs which they conjecture to be mirror to the anti-canonical hypersurfaces in G/P . In particular, they degenerate the pair $(G/B, -K_{G/B})$ to a toric pair $(X_\Delta, -K_{X_\Delta})$ corresponding to a certain polytope Δ . The polytope Δ has a dual polytope Δ^* corresponding to a toric pair $(X_{\Delta^*}, -K_{X_{\Delta^*}})$. They define a subfamily of $| -K_{X_{\Delta^*}} |$ whose coefficients satisfy "box equations". They conjecture that minimal crepant resolutions of generic elements of this subfamily are mirror to generic anti-canonical hypersurfaces in G/B . As evidence, they prove that the varieties have the same associated hypergeometric series [BCFKvS00, 5.2.1].

Following Caldero's work in [Cal02], Alexeev and Brion have constructed many different degenerations of spherical varieties [AB04]. These constructions include degenerations of the flag variety indexed by reduced decompositions $\overline{\omega}_0$ of ω_0 , the longest element in the Weyl group of G . In particular $(G/B, -K_{G/B})$ degenerates to a toric pair $(X_\Delta, \mathcal{O}_{X_\Delta}(1))$ corresponding to a rational polytope $\Delta = \Delta(\overline{\omega}_0)$ [AB04]. We call these degenerations *string degenerations*. The fibers in the limit vary greatly depending on the choice of reduced decomposition. For example, the polytope's F-vector (number of faces of different dimensions), the rank of the Picard group of X_Δ , and the dimension of the linear system $|\mathcal{O}_{X_{\Delta^*}}(1)|$ may all differ.

We would like to construct a mirror family using any of the string degenerations and see how they differ depending on the choice of degeneration. For a small toric degeneration (see Definition 5.1) Batyrev proposed such a family [Bat04].

Date: February 2, 2008.

We prove that $\Delta^*(\overline{\omega}_0)$ is integral, so it corresponds to a toric variety X_{Δ^*} with an ample line bundle $\mathcal{O}_{X_{\Delta^*}}(1)$. For any choice of reduced composition $\overline{\omega}_0$, we define $F_{\overline{\omega}_0}$ as a subfamily of $|\mathcal{O}_{X_{\Delta^*}}(1)|$ whose coefficients satisfy a set of relations we call *combinatorial box equations*. We prove that this family coincides with the family in Batyrev's construction when the string degeneration is small.

Using Berenstein and Zelevinsky's concept of geometric lifting [BZ01], we construct birational maps between any two families $F_{\overline{\omega}_0}$ and $F_{\overline{\omega}_0'}$. By the work of Batyrev and Kontsevich [Bat00, K95], this shows that smooth Calabi-Yau varieties birational to the generic elements of $F_{\overline{\omega}_0}$ all have the same Hodge numbers.

The organization of the paper is as follows:

In section 2 we fix notation and describe the string degenerations of G/B . We review Gleizer and Postnikov's description of the string cone [GP00] and use it to define a family of varieties $F_{\overline{\omega}_0}$.

Every pair of reduced decompositions of ω_0 is connected by a sequence of braid moves. In section 3, we classify how the inequalities defining the polytopes Δ change under a braid move. This allows us to prove that the polytopes Δ^* are integral, and will help us in proving that $F_{\overline{\omega}_0}$ and $F_{\overline{\omega}_0'}$ are birational.

In section 4, we define a birational map between the families $F_{\overline{\omega}_0}$ and $F_{\overline{\omega}_0'}$. This map is closely related to Berenstein and Zelevinsky's geometric lifting of piecewise linear maps to maps between totally positive varieties [BZ01].

In section 5, we prove that the families $F_{\overline{\omega}_0}$ coincide with the families that Batyrev constructed when the string degeneration is small. Finally, we give an example of a string degeneration which is not small.

Acknowledgments. The author would like to thank his advisor Valery Alexeev for posing this interesting problem, and for his helpful guidance and advice. The author would also like to thank Robert Varley for many helpful conversations.

2. PRELIMINARIES

We begin by introducing notation that will be used throughout this paper. Working over $\mathbb{C}$, let G be the algebraic group SL_{n+1} , B the Borel subgroup of upper triangular matrices, U the subgroup of upper uni-triangular matrices, and T the maximal torus of diagonal matrices. One has $B = TU$. Let Φ be the root system of (G, T) , with Φ^+ the subset of positive roots. We denote $\alpha_1, \dots, \alpha_n$ the corresponding simple roots.

Let W be the Weyl group of (G, T) with $s_1, \dots, s_n$ the reflections associated to $\alpha_1, \dots, \alpha_n$. Note that $W \cong S_{n+1}$ so we can view s_i as the transposition exchanging i and $i+1$. These transpositions define a length function l on W . Let ω_0 be the unique element of maximal length corresponding to the permutation sending 1 to $n+1$, 2 to n and so forth. There are $N = l(\omega_0) = \frac{n(n+1)}{2}$ positive roots, which is the dimension of the flag variety G/B .

We say that a decomposition $\overline{\omega} = s_{i_1} s_{i_2} \dots s_{i_m}$ of ω is reduced if $m = l(\omega)$. For every reduced decomposition of ω_0 , the string degeneration is a flat degeneration of $(G/B, -K_{G/B})$ to a toric variety X_{Δ} and a corresponding $\mathbb{Q}$ -line bundle $\mathcal{O}_{X_{\Delta}}(1)$ [AB04, Cal02]. By standard methods (see [Ful93, Bat94]) one can associate to $(X_{\Delta}, \mathcal{O}_{X_{\Delta}}(1))$ a polytope $\Delta = \Delta(\overline{\omega}_0)$ and its dual polytope $\Delta^* = \Delta^*(\overline{\omega}_0)$. For any reduced decomposition, Δ is a rational polytope in $\mathbb{R}^N$, thus X_{Δ} is a toric variety of dim N . For $n \leq 4$ and any choice of reduced decomposition, Alexeev and Brion verified that Δ is integral [AB04, Ex. 5.6]. This implies that the $\mathbb{Q}$ -line bundle

is actually a line bundle. In Section 3 we prove that for any n and any choice of reduced decomposition, Δ^* is an integral polytope and thus corresponds to a toric variety X_{Δ^*} with ample line bundle $\mathcal{O}_{X_{\Delta^*}}(1)$.


The polytope Δ is the intersection of Berenstein-Zelevinsky's string cone and a polyhedral cone called the λ -cone [AB04, Thm 1.1]. Explicit inequalities describing the λ -cone can be found in [Lit98, 1] and [AB04, Thm 1.1]. Inequalities for the string cone are given in [BZ01, 3.10] and [Lit98, 1]. A combinatorial description for them is given in [GP00, Cor.5.8].

Gleizer and Postnikov's description of the string cone is crucial to what follows so we review it here. (Note: in this review the order in which the s_i appear is reversed from that of the original.)

Let $\overline{\omega_0} = s_{i_1} s_{i_2} \dots s_{i_N}$ be a reduced decomposition of ω_0 . They represent this decomposition with a string diagram described below: Start with $n + 1$ strings at the top of the diagram. Move down the diagram exchanging the string in the i_1 column with the string in the $i_1 + 1$ column. Continuing down the diagram exchange the string in the i_2 column with the string in the $i_2 + 1$ column. Continue this process until you have made N exchanges. The strings will have reversed their orders when they reach the bottom of the diagram.

Label the intersection points $t_1, t_2, \dots, t_N$ from top to bottom. Label the strings U_1 through U_{n+1} on the top from left to right. Mark vertices u_1 through u_{n+1} (resp. b_1 through b_{n+1}) as the upper (resp. lower) ends of the strings U_1 through U_{n+1} .

Example 2.1. Here is a picture of the string diagram for $n = 3$ and reduced word decomposition $\overline{\omega_0} = s_3 s_1 s_2 s_1 s_3 s_2$:



Definition 2.2. ([GP00]) Pick a number k from 1 to n . Form an oriented graph on the string diagram as follows. Orient strings U_1 through U_k upward and the rest of the strings downward. A *rigorous path* is an oriented path from b_k to b_{k+1} not meeting any vertex more than once, and avoiding the two bad fragments pictured below.



Here the thin line denotes a string, and the thick line indicates the rigorous path.

Each rigorous path defines an inequality as follows:

$$\sum a_i x_i \geq 0 \text{ where } a_i = \begin{cases} 1 & \text{if the path switches from } U_i \text{ to } U_j \text{ and } i < j \\ -1 & \text{if the path switches from } U_i \text{ to } U_j \text{ and } i > j \end{cases}$$

We refer to these inequalities as string inequalities.

Example 2.3. For $k = 1$ in the string diagram pictured above, the rigorous path $b_1 \mapsto t_5 \mapsto t_3 \mapsto t_4 \mapsto t_6 \mapsto b_2$ corresponds to the inequality: $x_3 \geq x_4$.

Gleizer and Postnikov prove that the string cone is the collection of points in $\mathbb{R}^N$ satisfying the string inequalities [GP00, Cor. 5.8].

Lemma 2.4. *The λ -cone is the collection of points in $\mathbb{R}^N$ satisfying the following inequalities which we call λ -inequalities:*

$$\lambda_i : x_i \leq 2 + \sum_{j>i} c_j x_j$$

for i from 1 to N , where

$$c_j = \begin{cases} 1 & \text{if the vertex } t_j \text{ is one column to the right or left of } t_i \\ -2 & \text{if the vertex for } t_j \text{ is in the same column as } t_i \\ 0 & \text{otherwise.} \end{cases}$$

This is simply a way of visualising the Lie-algebraic definition of the λ -cone (see [AB04, Thm 1.1]), and of associating a λ -inequality to each intersection point in the string diagram.

Example 2.5. For the string diagram pictured above we have

$$\lambda_2 : x_2 \leq 2 + x_3 - 2x_4 + x_6.$$

Definition 2.6. Write the defining inequalities of $\Delta(\overline{\omega_0})$ as $\sum_i m_{i_d} x_i \leq (0 \text{ or } 2)$, and define $M_d = (m_{1_d}, \dots, m_{N_d})$. Let $T^{M_d} := t_1^{m_{1_d}} t_2^{m_{2_d}} \dots t_N^{m_{N_d}}$ and define $BF_{\overline{\omega_0}}$ as the family of hypersurfaces in the torus $T = \text{Spec} \mathbb{C}[t_1, t_1^{-1}, \dots, t_N, t_N^{-1}]$ given by the equations:

$$f_{\overline{\omega_0}, a} := 1 - \sum_d a_d T^{M_d} = 0$$

where d runs over all defining inequalities of Δ . We will refer to individual hypersurfaces as $Z_{\overline{\omega_0}, a}$, where $a = (a_1, \dots, a_r)$ denotes the coefficient vector.

Example 2.7. The λ_2 -inequality above would correspond to the term $a_{\lambda_2} t_2 t_3^{-1} t_4^2 t_6^{-1}$.

Note that $BF_{\overline{\omega_0}}$ is a family over the affine space $\mathbb{A}^r$. We will see in Section 3 that it is a subfamily of the linear system $|\mathcal{O}_{X_{\Delta^*}}(1)|$ intersected with the torus.

We restrict our attention to a subfamily whose coefficients satisfy a set of relations we call *combinatorial box equations*. We define these relations in terms of the bounded regions of the string diagram which we call *boxes*. Every box of the string diagram is obviously bounded above by a vertex t_{top} and directly below by a vertex t_{bot} . Let $T^{\lambda_{\text{top}}}$ (resp. $T^{\lambda_{\text{bot}}}$) be the monomial in $f_{\overline{\omega_0}, a}$ corresponding to the λ -inequality associated to t_{top} (resp. t_{bot}).

Definition 2.8. For every two string inequalities p_1, p_2 with corresponding monomials T^{p_1} and T^{p_2} satisfying the following *box conditions*:

- (1) there exists a box with corresponding monomials $T^{\lambda_{\text{top}}}$ and $T^{\lambda_{\text{bot}}}$ such that $T^{p_1} T^{\lambda_{\text{top}}} = T^{p_2} T^{\lambda_{\text{bot}}}$,
- (2) the t_{top} degree of $T^{p_1} = -1$,
- (3) the t_{bot} degree of $T^{p_2} = 1$,

we define an equation $a_{p_1} a_{\lambda_{\text{top}}} = a_{p_2} a_{\lambda_{\text{bot}}}$. We call the collection of all such equations the *combinatorial box equations*.

Definition 2.9. Let $P_{\overline{\omega_0}}$ be the subscheme defined by the combinatorial box equations in the torus $\text{Spec } \mathbb{C}[a_1, a_1^{-1}, \dots, a_r, a_r^{-1}]$.

Definition 2.10. Let $F_{\overline{\omega_0}}$ be the subfamily of $BF_{\overline{\omega_0}}$ whose coefficients are non-zero and satisfy the combinatorial box equations. Note that $F_{\overline{\omega_0}}$ is a family over the base $P_{\overline{\omega_0}}$.

It is a standard fact that any two reduced decompositions of ω_0 are connected by a finite sequence of the following two braid moves.

- (1) 2-move: exchanges (s_i, s_j) with (s_j, s_i) where $|i - j| > 1$
- (2) 3-move: exchanges (s_i, s_j, s_i) with (s_j, s_i, s_j) where $|i - j| = 1$

A 3-move from $\overline{\omega_0}$ to $\overline{\omega'_0}$ fixes all of the string diagram except for one box.

Definition 2.11. Given a 3-move from $\overline{\omega_0}$ to $\overline{\omega'_0}$, label the box in the string diagram for $\overline{\omega_0}$ bounded by t_i, t_j and t_k , R and its image R' . We say that R is of type $R121$ if it is of the form (s_i, s_{i+1}, s_i) . Otherwise we say it is of type $R212$. We refer to these regions as braid regions. Note that if R is of type $R121$ then R' must be of type $R212$ and vice versa.

The map between string cones differing by a single braid move, defined by Berenstein and Zelevinsky in [BZ93, Thm 2.7], restricts to a map between polytopes $\Delta(\overline{\omega_0})$ and $\Delta(\overline{\omega'_0})$.

They prove that the map fixes all of the coordinates except for the two or three corresponding to those being exchanged in the braid move. For those coordinates they have the following maps:

- (1) 2-move: $(x_i, x_j) \rightarrow (x_j, x_i)$
- (2) 3-move: $(x_i, x_j, x_k) \rightarrow (\max(x_k, x_j - x_i), x_i + x_k, \min(x_i, x_j - x_k))$.

3. CLASSIFICATION OF HOW Δ CHANGES UNDER 3-MOVE

In this section we classify the defining inequalities of Δ and how those inequalities change under a 3-move. From these results we prove that the dual polytope Δ^* is integral.

Only three coordinates change under a 3-move. In what follows we will refer to those coordinates as x_i, x_j , and x_k (note that i, j , and k are consecutive integers). We call the product of the t_i, t_j , and t_k variables inside of T^{M_d} a *monomial piece*, and refer to this product as τ^{M_d} . We will refer to the monomial pieces associated to string (resp. λ) inequalities as string (resp. λ) monomial pieces.

Since each inequality corresponds to a monomial, we can classify all possible string and λ -inequalities in terms of their monomial pieces.

3.1. Classification of λ -Monomial Pieces.

Theorem 3.1. *The following is a complete classification of λ -monomial pieces before and after a 3-move.*


$$\begin{array}{lllll} t_i t_j^{-1} t_k^2 \leftrightarrow t_i t_j^{-1} t_k^2 & t_j t_k^{-1} \leftrightarrow t_j t_k^{-1} & t_k \leftrightarrow t_k & t_i^{-1} t_k^{-1} \leftrightarrow t_j^{-1} \\ t_i^2 t_j^{-1} t_k^2 \leftrightarrow t_i^{-1} t_j^2 t_k^{-1} & & & \end{array}$$

An arrow indicates which monomial pieces are exchanges under the braid move.

Proof. As above, we fix t_i, t_j and t_k as the coordinates on which the 3-move occurs. We classify the monomial pieces for the inequalities λ_l for all values of l . For $l > k$ the monomial for λ_l doesn't have any terms from the braid move section. This means that the monomial piece is 1 and it doesn't change after a braid move.

For λ_i, λ_j , and λ_k we have the following monomial pieces regardless if R is of type $R121$ or $R212$: $\lambda_i : t_i t_j^{-1} t_k^2$ $\lambda_j : t_j t_k^{-1}$ $\lambda_k : t_k$

If $l < i$, then the λ_l monomial piece only depends on which column t_l lies (as pictured below).



- (1) If t_l is in column 1 we have the monomial piece $t_i^{-1} t_k^{-1}$ before, and t_j^{-1} after the braid move.
- (2) If t_l is in column 2 we have the monomial piece $t_i^2 t_j^{-1} t_k^2$ before, and $t_i^{-1} t_j^2 t_k^{-1}$ after the braid move.
- (3) If t_l is in column 3 we have the monomial piece $t_i^{-1} t_j^2 t_k^{-1}$ before, and $t_i^2 t_j^{-1} t_k^2$ after the braid move.
- (4) If t_l is in column 4 we have the monomial piece t_j^{-1} before, and $t_i^{-1} t_k^{-1}$ after the braid move.

Note that for R of type $R212$ we get the same collection of monomial pieces. $\square$

3.2. Classification of String Monomial Pieces.


Theorem 3.2. *The following is a complete classification of the string monomial pieces, and how they are exchanged under a type B_2 braid move. The $\&$ sign indicates that a pair of monomial pieces must occur together.*

$$\begin{aligned}
 t_k^{-1} &\leftrightarrow t_i^{-1} \& t_j^{-1} t_k \\
 t_i &\leftrightarrow t_k \& t_i^{-1} t_j \\
 t_i t_k &\leftrightarrow t_j \\
 t_i^{-1} t_k^{-1} &\leftrightarrow t_j^{-1} \\
 t_i t_j^{-1} t_k &\leftrightarrow t_i^{-1} t_j t_k^{-1}
 \end{aligned}$$


Proof. We would like to classify all of the string inequalities in terms of their monomial pieces. By symmetry we can assume that R is of type $R121$. If a string monomial piece is anything other than 1, the corresponding rigorous path p must change strings in the region R . Any oriented path entering R more than once must intersect itself so it cannot be rigorous. Therefore, we may assume that a rigorous path only enters and exits R once. There are $6 * 5 = 30$ possible pairs of entry and exit points for p . If p enters R at the vertex b_e , then all strings U_f for $f \geq e$ must be oriented upward. Therefore p cannot exit R through b_f where $f < e$. Similarly if p enters at u_e it may not exit through u_f for $f > e$. This eliminates 6 pairs

of entry and exit points. By the same reasoning no rigorous path which changes strings on R can enter at b_3 (resp. u_1) and exit at u_3 (resp. b_1). What remains are 22 entry and exit pairs.



The following three diagrams correspond to cases of entry and exit points in which there is a unique rigorous path before and after the braid move.



In the following three diagrams there is one possible rigorous path through R before the braid move. After the braid move there are two possible rigorous paths through R' . Since the string cone is the collection of points satisfying all possible string inequalities there must be two inequalities (and hence monomial pieces) in the image. In what follows we group these monomial pieces together.



We have the monomial piece $t_k^{-1} \rightarrow t_i^{-1}$ & $t_j^{-1} t'_k$



We have the monomial pieces $t_k^{-1} \rightarrow t_i^{-1}$ & $t_j^{-1} t'_k$.

We can construct the other 16 cases of entry and exit points from these six using symmetry. For each rigorous path on R' listed above there is a corresponding rigorous path on R as follows. For the entry and exit points of the new path, exchange b_1 and b_3 (resp u_1 and u_3). Then construct a new path which changes strings at the same vertices as the path on R' . Unless the path on R' changes strings at every vertex, this new path will be rigorous. This is what we call horizontal path symmetry. Rigorous paths which changes strings at every vertex do not have horizontally symmetric partners. There is also a vertical symmetry. Take a path on R and exchange the “top” with “bottom” on the entry and exit points (i.e. b_3 is switched with u_1). Now create a path which changes strings at vertically flipped vertices (i.e. t_i instead of t_k). This will be a new rigorous path. The remaining 16 single entry point cases can be gotten from the 6 listed above, through a combination of horizontal and vertical symmetries. From this classification we see that the sets of monomial pieces stated in the theorem are exchanged under a 3-move. $\square$

3.3. Integrality of Δ^* .

Lemma 3.3. *For any braid region R , the line connecting the apex of the λ -cone and the origin lies on the hyperplane $x_i + x_k = x_j$.*

Proof. It suffices to show that the apex of the λ -cone lies on the hyperplane. At the apex all of the λ -inequalities are equalities. We can write the following equations, where A , B and are the contributions to the equation from x_d for $d > k$.

$$\begin{aligned} x_i &= 2 + x_j - 2x_k + A \\ x_k &= 2 + B \end{aligned}$$

Note that since t_i and t_k are in the same column of the string diagram we have $A = B$. From this we can see that

$$x_i + x_k = (2 + x_j - 2x_k + A) + x_k = (x_j - x_k) + 2 + A = (x_j - x_k) + x_k = x_j.$$

$\square$

Theorem 3.4. $\Delta^*(\overline{\omega_0})$ is an integral polytope for any reduced decomposition $\overline{\omega_0}$.

Proof. In [BCFKvS00], Batyrev (et. al.) show that for the reduced decomposition $\overline{\omega_0} = s_1 s_2 s_1 \dots s_n s_{n-1} \dots s_1$ the polytope Δ^* is an integral polytope with a unique interior point $P = (1, 2, 1, \dots, n, n-1, \dots, 2, 1)$. The facets of Δ determine linear functions L such that $L(P) = 1$. Let Ψ be the piecewise linear map between polytopes defined by Berenstein and Zelevinsky. We want to show that under a 3-move, the facets of Δ' determine L' such that $L'(P') = 1$ where $P' = \Psi(P)$.

The map Ψ is defined by two linear maps ψ_1 and ψ_2 which agree on the hyperplane $x_i + x_k = x_j$. For any linear function L associated to a facet of Δ there are two corresponding linear functions on Δ' ; $L'_1 = L \circ \psi_1^{-1}$ and $L'_2 = L \circ \psi_2^{-1}$.

The point P is halfway between the origin (which is the apex of the string cone) and the apex of the λ -cone. By the preceding lemma this shows that P lies on the hyperplane $x_i + x_k = x_j$. Since ψ_1 and ψ_2 agree on that hyperplane we have $\psi_1(P) = \psi_2(P) = \Psi(P)$.

Therefore, we can make the calculations $L'_1(P') = L(\psi_1^{-1}(P')) = L(P) = 1$ and $L'_2(P') = L(\psi_1^{-1}(P')) = L(P) = 1$, which implies that the polytope Δ'^* is integral. Note that P' is still halfway between the origin and the apex of the λ -cone, which allows this argument to be repeated.

Under a 2-move the polytopes Δ and Δ' are isomorphic, so Δ^* is integral, if and only if Δ'^* is integral.

By starting with $\overline{\omega}_0 = s_1 s_2 s_1 \dots s_n s_{n-1} \dots s_1$ and the point $P = (1, 2, 1, \dots, n, n-1, \dots, 2, 1)$ we can compose braid moves and always have an interior point P' on which all of the defining linear functionals take value 1. This proves that Δ^* is integral for any choice of reduced decomposition. $\square$

Corollary 3.5. *The monomials in $f_{\overline{\omega}_0, a}$ correspond to vertices of Δ^* and, $F_{\overline{\omega}_0}$ is a subfamily of the linear system $|\mathcal{O}_{X_{\Delta^*}}(1)|$ intersected with the torus.*

4. BIRATIONALITY OF $F_{\overline{\omega}_0}$

Definition 4.1. Assume that $\overline{\omega}_0$ and $\overline{\omega}_0'$ differ by a 3-move. Let $C = \frac{a_{\lambda_k}}{a_{\lambda_i}}$, $T = \text{Spec } \mathbb{C}[t_1, t_1^{-1}, \dots, t_N, t_N^{-1}]$ and $T' = \text{Spec } \mathbb{C}[t'_1, t'_1{}^{-1}, \dots, t'_N, t'_N{}^{-1}]$. For a fixed coefficient vector a , define $h_a : T' \dashrightarrow T$ by $(t_i, t_j, t_k) = (\frac{t'_i t'_k + C t'_j}{t'_i}, \frac{t'_i t'_k}{C}, \frac{C t'_i t'_j}{t'_i t'_k + C t'_j})$, and $t_q = t'_q$ for $q \notin \{i, j, k\}$.

Define a map $h'_a : T \dashrightarrow T'$ by exchanging t with t' in the map above. Then we have $h'_a \circ h_a = \text{id}$.

Proposition 4.2. *For a fixed coefficient vector $a = (a_1, \dots, a_r) \in P_{\overline{\omega}_0}$, $f'_a := h_a^*(f_{\overline{\omega}_0, a})$ defines a variety $Z_{\overline{\omega}_0', a'}$ for some coefficient vector a' . Therefore, we have an induced birational map $h_a : Z_{\overline{\omega}_0', a'} \dashrightarrow Z_{\overline{\omega}_0, a}$.*

Remark 4.3. We will see in Proposition 4.9 that the map g defined by $g^*(a) = a'$ is an isomorphism between the parameter spaces $P'_{\overline{\omega}_0'}$ and $P_{\overline{\omega}_0}$. The coordinates for a' are explicitly given as regular functions of the coordinates for a in Proposition 4.6.

Remark 4.4. The map h_a is a specific geometric lift (in the sense of [BZ01]) of the piecewise linear function: $(t'_i, t'_j, t'_k) \rightarrow (\min(t'_k, t'_j - t'_i), t'_i + t'_k, \max(t'_i, t'_j - t'_k))$.

Proof of Proposition 4.2. We show that for a fixed coefficient vector a , h_a is birational in two steps. First, in Proposition 4.6, we show that $h_a^*(1 - \sum_d a_d T^{M_d}) = (1 - \sum_{d'} a'_d T^{M'_d})$ for some coefficient vector a' . Then, in Section 4.2 we show that the coefficient vector a' is actually in $P'_{\overline{\omega}_0'}$ by showing that g^* preserves the combinatorial box equations.

4.1. Action of h^* on $f_{\overline{\omega}_0, a}$.

Definition 4.5. We refer to the box equations coming from the boxes R and R' as *local box equations*.

Proposition 4.6. *Fix a coefficient vector a satisfying the local box equations. Then $h_a^*(1 - \sum_d a_d T^{M_d}) = (1 - \sum_{d'} a'_d T^{M'_d})$ for some coefficient vector a' .*

Proof. We examine how h_a^* acts on $1 - \sum_d a_d T^{M_d}$ monomial by monomial. Since h_a^* only depends on t_i, t_j , and t_k we classify its action on monomials based on the corresponding monomial pieces.

A priori $h_a^*(f_{\overline{\omega}_0, a})$ is a rational function, but not a sum of monomials. In several cases, in order to see that $h_a^*(1 - \sum_d a_d T^{M_d})$ is a linear combination of monomials, we have to group pairs of monomials together. In some cases h_a^* only maps a particular combination of two monomials to a monomial, if the coefficients satisfy the local box equations.

The map h_a^* take the following classes of monomials to monomials (or sum of monomials):

$$\begin{aligned}
h_a^*(a_d t_i^{-1} t_k^{-1}) &= a_d a_{\lambda_i} a_{\lambda_k}^{-1} t_j'^{-1} \\
h_a^*(a_d t_i^2 t_j^{-1} t_k^2) &= a_d a_{\lambda_i}^{-3} a_{\lambda_k}^3 t_i'^{-1} t_j'^2 t_k'^{-1} \\
h_a^*(a_d t_i^{-1} t_j^2 t_k^{-1}) &= a_d a_{\lambda_i}^3 a_{\lambda_k}^{-3} t_i'^2 t_j'^{-1} t_k'^2 \\
h_a^*(a_d t_j^{-1}) &= a_d a_{\lambda_i}^{-1} a_{\lambda_k} t_i'^{-1} t_k'^{-1} \\
h_a^*(a_d t_i^{-1} t_k^{-1}) &= a_d a_{\lambda_i} a_{\lambda_k}^{-1} t_j'^{-1} \\
h_a^*(a_d t_j^{-1}) &= a_d a_{\lambda_i}^{-1} a_{\lambda_k} t_i'^{-1} t_k'^{-1} \\
h_a^*(a_d t_i t_k) &= a_d a_{\lambda_i}^{-1} a_{\lambda_k} t_j' \\
h_a^*(a_d t_j) &= a_d a_{\lambda_i} a_{\lambda_k}^{-1} t_i' t_k' \\
h_a^*(a_d t_i t_j^{-1} t_k) &= a_d a_{\lambda_i}^2 a_{\lambda_k}^{-2} t_i'^{-1} t_j' t_k'^{-1} \\
h_a^*(a_{\lambda_j} t_j t_k^{-1}) &= a_{\lambda_i}^2 a_{\lambda_j} a_{\lambda_k}^{-2} t_i'^{-1} t_j'^2 t_k'^{-1} + a_{\lambda_i} a_{\lambda_j} a_{\lambda_k}^{-1} t_k' \\
h_a^*(a_d t_i^{-1} t_j t_k^{-1}) &= a_d a_{\lambda_i}^{-2} a_{\lambda_k}^2 t_i' t_j'^{-1} t_k' \\
h_a^*(a_d t_k^{-1}) &= a_d t_i'^{-1} + a_d a_{\lambda_i} a_{\lambda_k}^{-1} t_j'^{-1} t_k' \\
h_a^*(a_d t_i) &= a_d t_k' + a_d a_{\lambda_i}^{-1} a_{\lambda_k} t_i'^{-1} t_j'
\end{aligned}$$

The following classes of monomials must be grouped together in order for h_a^* to take them to a monomial.

$$h_a^*(a_{\lambda_i} t_i t_j^{-1} t_k^2 + a_{\lambda_k} t_k) = a_{\lambda_i}^{-1} a_{\lambda_k}^2 t_j' t_k'^{-1}$$

The following classes of monomials must be grouped together, and their coefficients must satisfy the local box equations in order for h_a^* to take them to monomials. We write the second coefficient in terms of the local box equation.

$$\begin{aligned}
h_a^*(a_d t_k + a_d a_{\lambda_i}^{-1} a_{\lambda_k} t_i^{-1} t_j) &= a_d t_i' \\
h_a^*(a_d t_i^{-1} + a_d a_{\lambda_i} a_{\lambda_k}^{-1} t_j^{-1} t_k) &= a_d t_k'^{-1}
\end{aligned}$$

The monomials occurring in $\text{image}(h_a^*)$ are precisely those occurring in $f_{\overline{\omega}_0, a}$. From this we see that

$h_a^*(1 - \sum_d a_d T^{M_d}) = (1 - \sum_{d'} a'_d T^{M'_d})$ for some coefficient vector a' . The inverse map is given by using the same construction for the braid move from $\overline{\omega}_0'$ to $\overline{\omega}_0$. $\square$

4.2. Preservation of Box Equations.

Proposition 4.7. *For any 3-move, the map g^* preserves box equations.*

Proof. Let R' be the image of the box R (We can assume it is of type $R212$ with the proof of the other case following using the same methods). We classify all boxes O' of the string diagram for $\overline{\omega}_0'$ by their positions in comparison with R' .

The proof of every case follows the following method: First, classify all possibilities of monomial pieces τ for which the box condition on O' could be satisfied. Since we know how the monomial pieces change under the braid move, we verify that the box conditions must be satisfied for the box $O = \text{preimage}(O')$. Since $a \in P$ we know that if the box conditions are satisfied on O , then the corresponding box equations (in a_d) are also satisfied. We use the map g to write a'_d in terms of a_d , and check that new combinatorial box equations on O' are satisfied.

Remark 4.8. For some string monomials on O' there are two different corresponding string monomials in the preimage. Only one of these two string monomials is used to construct a box condition on the preimage. The first box condition may not be met in the preimage when the other choice of string monomial is used.

We classify the boxes O' into eight groups. For each group, we state how many pair of monomial pieces could possibly correspond to monomials satisfying the box conditions on O' .

- (1) O' doesn't touch R' (1 pair)
- (2) $O' = R'$ (1 pair)
- (3) O' is below and left of R' (3 pair)
- (4) O' is above and left of R' (3 pair)
- (5) O' is directly above R' (3 pair)
- (6) O' is directly below R' (3 pair)
- (7) O' is to the right of R' (6 pair)
- (8) O' is to the left of R (6 pair)

What follows is the proof for the case when R' is directly above O' . The proof for all of the other groups follow the exact same method.

Assume R' is directly above O' as pictured below.



Assume that p'_1 and p'_2 satisfy the box conditions on O' . Let τ^{M_d} be the monomial piece associated to T^{M_d} . By comparing the λ -inequalities see the following

relationship among monomial pieces.

$$\frac{\tau^{p'_2}}{\tau^{p'_1}} = \frac{\tau^{\lambda_{\text{top}'}}}{\tau^{\lambda_{\text{bot}'}}} = t_k.$$

From this we see that there are three possibilities for pairs string monomial pieces.

(1) Assume that

$$\tau^{p'_1} = t_i^{-1} t_k^{-1} \text{ and } \tau^{p'_2} = t_i^{-1}.$$

We construct a set of monomials satisfying the box conditions for the box O as follows.

$$\begin{aligned} T^{p'_1} \cdot T^{\lambda_{\text{top}'}} &= T^{p'_2} \cdot T^{\lambda_{\text{bot}'}} \\ T^{p'_1} \cdot T^{\lambda_{\text{top}}} \cdot t_j^{-1} t_k^2 &= T^{p'_2} \cdot T^{\lambda_{\text{bot}}} \\ T^{p_1} \cdot t_i^{-1} t_j t_k^{-1} \cdot T^{\lambda_{\text{top}}} \cdot t_j^{-1} t_k^2 &= T^{p_2} \cdot t_i^{-1} t_k \cdot T^{\lambda_{\text{bot}}} \\ T^{p_1} \cdot T^{\lambda_{\text{top}}} &= T^{p_2} \cdot T^{\lambda_{\text{bot}}} \end{aligned}$$

Since the box condition are satisfied on O , we have the following equations in a_i .

$$a_{p_1} a_{\lambda_{\text{top}}} = a_{p_2} a_{\lambda_{\text{bot}}}.$$

Rewriting the a'_i in terms of a_i we see that

$$a_{p'_1} a_{\lambda_{\text{top}'}} = a_{p_1} a_{\lambda_{\text{top}}} \text{ and } a_{p'_2} a_{\lambda_{\text{bot}'}} = a_{p_2} a_{\lambda_{\text{bot}}}.$$

Therefore,

$$a_{p'_1} a_{\lambda_{\text{top}'}} = a_{p'_2} a_{\lambda_{\text{bot}'}}.$$

(2) Assume that

$$\tau^{p'_1} = t_k^{-1} \text{ and } \tau^{p'_2} = 1.$$

We construct a set of monomials satisfying the box conditions for the box O as follows.

$$\begin{aligned} T^{p'_1} \cdot T^{\lambda_{\text{top}'}} &= T^{p'_2} \cdot T^{\lambda_{\text{bot}'}} \\ T^{p'_1} \cdot T^{\lambda_{\text{top}}} \cdot t_j^{-1} t_k^2 &= T^{p'_2} \cdot T^{\lambda_{\text{bot}}} \\ T^{p_1} \cdot t_j t_k^{-2} \cdot T^{\lambda_{\text{top}}} \cdot t_j^{-1} t_k^2 &= T^{p_2} \cdot T^{\lambda_{\text{bot}}} \\ T^{p_1} \cdot T^{\lambda_{\text{top}}} &= T^{p_2} \cdot T^{\lambda_{\text{bot}}} \end{aligned}$$

Since the box condition were satisfied on O , we have the following equations in a_i .

$$a_{p_1} a_{\lambda_{\text{top}}} = a_{p_2} a_{\lambda_{\text{bot}}}.$$

Rewriting the a'_i in terms of a_i we see that

$$a_{p'_1} a_{\lambda_{\text{top}'}} = a_{p_1} a_{\lambda_{\text{top}}} \text{ and } a_{p'_2} a_{\lambda_{\text{bot}'}} = a_{p_2} a_{\lambda_{\text{bot}}}.$$

Therefore,

$$a_{p'_1} a_{\lambda_{\text{top}'}} = a_{p'_2} a_{\lambda_{\text{bot}'}}.$$

(3) Assume that

$$\tau^{p'_1} = t_i^{-1} t_j t_k^{-1} \text{ and } \tau^{p'_2} = t_i^{-1} t_j.$$

We construct a set of monomials satisfying the box conditions for the box O as follows.

$$\begin{aligned} T^{p'_1} \cdot T^{\lambda_{\text{top}'}} &= T^{p'_2} \cdot T^{\lambda_{\text{bot}'}} \\ T^{p'_1} \cdot T^{\lambda_{\text{top}}} \cdot t_j^{-1} t_k^2 &= T^{p'_2} \cdot T^{\lambda_{\text{bot}}} \\ T^{p_1} \cdot t_i^{-2} t_j \cdot T^{\lambda_{\text{top}}} \cdot t_j^{-1} t_k^2 &= T^{p_2} t_i^2 t_j^{-2} t_k^2 \cdot T^{\lambda_{\text{bot}}} \\ T^{p_1} \cdot T^{\lambda_{\text{top}}} &= T^{p_2} \cdot T^{\lambda_{\text{bot}}} \end{aligned}$$

Since the box condition are satisfied on O , we have the following equations in a_i .

$$a_{p_1} a_{\lambda_{\text{top}}} = a_{p_2} a_{\lambda_{\text{bot}}}.$$

Rewriting the a'_i in terms of a_i we see that

$$a_{p'_1} a_{\lambda_{\text{top}'}} = a_{p_1} \frac{a_k^2}{a_i^2} a_{\lambda_{\text{top}}} \frac{a_i}{a_k} \text{ and } a_{p'_2} a_{\lambda_{\text{bot}'}} = a_{p_2} \frac{a_k}{a_i} a_{\lambda_{\text{bot}}}.$$

Therefore,

$$a_{p'_1} a_{\lambda_{\text{top}'}} = a_{p'_2} a_{\lambda_{\text{bot}'}}.$$

We have shown that the combinatorial box equations are preserved under g^* , which completes the proof of proposition 4.2. $\square$

4.3. Isomorphism of Parameter Spaces.

Proposition 4.9. *If $\overline{\omega_0}$ and $\overline{\omega'_0}$ differ by a single braid move then $P'_{\overline{\omega_0}} \cong P_{\overline{\omega_0}}$.*

Proof. Assume $\overline{\omega_0}$ and $\overline{\omega'_0}$ differ by a 3-move. As in remark 4.3, define $g : P' \rightarrow P$ by $g^*(a) = a'$. By the proof of Proposition 4.6 we see that this defines a regular map from $P_{\overline{\omega_0}}$ to the torus of coordinates of $P'_{\overline{\omega'_0}}$. Proposition 4.7 shows that the image of this map is actually in $P'_{\overline{\omega_0}}$. Similarly we can construct a regular map $g' : P_{\overline{\omega_0}} \rightarrow P'_{\overline{\omega'_0}}$. We can use Proposition 4.6 to verify that these maps are inverse to one another.

To complete the proof we note that $P_{\overline{\omega_0}}$ and $P'_{\overline{\omega'_0}}$ are obviously isomorphic under a 2-move. $\square$

Corollary 4.10. *The parameter space $P_{\overline{\omega_0}}$ is a toric variety for any choice of $\overline{\omega_0}$.*

Proof. In the case of the standard reduced decomposition $(s_1 s_2 s_1 \dots s_n s_{n-1} \dots s_1)$ the combinatorial box equations and Batyrev's box equations are exactly the same. By Batyrev's work [Bat04, Rm.4.2], the set of non-zero a_i satisfying the box equations form a toric variety. By the previous lemma, we see that under a braid move the parameter spaces are isomorphic. We can compose braid moves to see that the parameter spaces are isomorphic for any choice of reduced decomposition. $\square$

4.4. Birationality of the Families $F_{\overline{\omega}_0}$.

Lemma 4.11. *For $\overline{\omega}_0$ and $\overline{\omega}_0'$ differing by a 2-move, the map $H : F_{\overline{\omega}_0'} \rightarrow F_{\overline{\omega}_0}$ defined by exchanging t_i with t_j and exchanging the corresponding coefficients is an isomorphism.*

Proof. This is a consequence of Berenstein-Zelevinsky's map between string cones for the 2-move case. $\square$

Theorem 4.12. *If $\overline{\omega}_0$ and $\overline{\omega}_0'$ differ by a 3-move, the map*

$$H : F_{\overline{\omega}_0'} \dashrightarrow F_{\overline{\omega}_0}$$

defined as $H := (g, h_a)$ is a birational map of families.

Proof. This follows from the fact that g is an isomorphism, and h_a is a birational map for any fixed coefficient vector a . $\square$

Corollary 4.13. *For any two reduced decompositions $\overline{\omega}_0$ and $\overline{\omega}_0'$, the families $F_{\overline{\omega}_0}$ and $F_{\overline{\omega}_0'}$ are birational.*

Proof. Connect $\overline{\omega}_0$ and $\overline{\omega}_0'$ by a sequence of braid moves. For each braid move there is a birational map of families, so we can compose these maps to get a birational map between $F_{\overline{\omega}_0}$ and $F_{\overline{\omega}_0'}$. $\square$

Corollary 4.14. *Smooth Calabi-Yau manifolds birational to the generic elements of $F_{\overline{\omega}_0}$ have the same Hodge numbers for any choice of $\overline{\omega}_0$.*

Proof. Since the generic elements are all birational, smooth Calabi-Yau manifolds birational to them have the same Hodge numbers by the work of Batyrev and Kontsevich [Bat00, K95]. $\square$

5. COMPARING $F_{\overline{\omega}_0}$ WITH THE FAMILIES DEFINED BY BATYREV (ET.AL.)

Definition 5.1. [Bat04, Def 3.1] Let $X \subset \mathbb{P}^m$ be a smooth Fano variety of dimension n . A normal Gorenstein toric Fano variety $Y \subset \mathbb{P}^m$ is called a *small toric degeneration* of X , if there exists a Zariski open neighbourhood U of $0 \in \mathbb{A}^1$ and an irreducible subvariety $\tilde{X} \subset \mathbb{P}^m \times U$ such that the morphism $\pi : \tilde{X} \rightarrow U$ is flat and the following conditions hold:

- (1) the fiber $X_t := \pi^{-1}(t) \subset \mathbb{P}^m$ is smooth for all $t \in U \setminus 0$;
- (2) the special fiber $X_0 := \pi^{-1}(0) \subset \mathbb{P}^m$ has at worst Gorenstein terminal singularities and X_0 is isomorphic to $Y \subset \mathbb{P}^m$;
- (3) the canonical homomorphism

$$\text{Pic}(\tilde{X}/U) \rightarrow \text{Pic}(X_t)$$

is an isomorphism for all $t \in U$.

In [Bat04] Batyrev proposes a mirror construction for any small toric degeneration. This construction is an extension of Batyrev (et. al.)'s construction in the case of partial flag varieties. We set up the next lemma to help show that our construction is the same as Batyrev's in the case of a small degeneration.

Lemma 5.2. *For any inequalities λ_{top} , λ_{bot} , p_1 , and p_2 whose corresponding monomials satisfy a combinatorial box condition on O , there exists a facet of Δ^* containing the vertices corresponding to λ_{top} , λ_{bot} , p_1 , and p_2 .*

Proof. Define the linear function $L(t_1, \dots, t_N) := t_{\text{top}} - t_{\text{bot}} + 1$. We check that $L = 0$ on the corresponding vertices $\lambda_{\text{top}}, \lambda_{\text{bot}}, p_1$, and p_2 , and $L \geq 0$ on all other vertices of Δ^* .

$$L(\lambda_{\text{top}}) = 1 - 2 + 1 = 0$$

$$L(\lambda_{\text{bot}}) = 0 - 1 + 1 = 0$$

$$L(p_1) = -1 - x + 1$$

$$L(p_2) = y - 1 + 1$$

But since $\lambda_{\text{top}} + p_1 = \lambda_{\text{bot}} + p_2$ we see that $x = y = 0$ which implies that $L(p_1) = L(p_2) = 0$.

Next we want to show that for every other vertex v of Δ^* , $L(v) \geq 0$. Assume v is a vertex corresponding to a λ -inequality λ_v then

$$L(v) = \begin{cases} 0 - 0 + 1 & = 1 \text{ If } \lambda_v \text{ lies below } \lambda_{\text{bot}} \text{ or least 2 columns to the} \\ & \text{left or right of } \lambda_{\text{top}} \\ 0 - (-1) + 1 & = 2 \text{ If } \lambda_v \text{ is in the region O.} \\ 2 - 2 + 1 & = 1 \text{ If } \lambda_v \text{ lies in the column above } \lambda_{\text{top}} \\ -1 - (-1) + 1 & = 1 \text{ If } \lambda_v \text{ lies above and adjacent to } \lambda_{\text{top}}. \end{cases}$$

String vertices can only take values from the set $\{-1, 0, 1\}$ on the coordinates t_{top} and t_{bot} . From this we see that the only way the linear function L could be negative on such a vertex v , is if t_{top} coordinate of v was -1 and the t_{bot} coordinate was 1 . Any such inequality would correspond to an oriented path which intersects itself, which is a contradiction.

We see that $(L = 0)$ defines a face of Δ containing the four vertices. This completes the proof since any face is contained in a facet. $\square$

Theorem 5.3. *For a small string degeneration $F_{\overline{\omega}_0}$ is same as the family constructed by Batyrev in [Bat04, Sect. 4].*

Proof. In the case of a small toric degeneration Batyrev defines a subfamily of varieties in $|\mathcal{O}_{X_{\Delta^*}}(1)|$ whose coefficients satisfy box equations. (Box equation terminology was used in [BCFKvS00]. The coefficients satisfying the box equations are referred to as Σ -admissible in [Bat04].)

Our combinatorial box equations are a subset of Batyrev's box equations if, and only if, the vertices in Δ^* corresponding to $\lambda_{\text{top}}, \lambda_{\text{bot}}, p_1$, and p_2 all lie in a common facet of Δ^* . Therefore, by Lemma 5.2 we have shown that the combinatorial box equations are a subset of Batyrev's box equations. Batyrev proves in [Bat04, Rem. 4.2] that his box equations define an irreducible variety of $\dim = \text{rank}(\text{Pic}(X_{\Delta^*})) + \dim(G/B)$. This is the same as the dimension of the parameter space $P_{\overline{\omega}_0}$. By Lemma 4.10, $P_{\overline{\omega}_0}$ is an irreducible variety. If Batyrev's family had any more independent box equations they would define a variety of smaller dimension. Therefore, Batyrev's box equations cut out exactly $P_{\overline{\omega}_0}$, and the families are the same. $\square$

Corollary 5.4. *For all reduced decompositions $\overline{\omega}_0$, generic elements of $F_{\overline{\omega}_0}$ share the mirror properties of the generic elements of Batyrev's families.*

Proof. In the small case the families are the same by Theorem 5.3, and thus their generic elements share the same mirror properties. By Corollary 4.12 the generic elements are birational for any choice of reduced decomposition, and therefore share the same mirror properties. $\square$

Remark 5.5. For the reduced decomposition $\overline{\omega}_0 = s_3 s_1 s_2 s_1 s_3 s_2$ calculation shows that $\text{rank}(\text{Pic}(X_\Delta)) < \text{rank}(\text{Pic}(G/B))$. Therefore, the corresponding string degenerations is not small. This calculation was performed with the aid of the freely available program `polymake` using the techniques found in [Ful93].

REFERENCES

- [AB04] V. Alexeev and M. Brion, *Toric Degenerations of Spherical Varieties*, Selecta Math. (N.S.) **10**(2004) no.4, 453-478.
- [Bat04] V.V. Batyrev, *Toric Degenerations of Fano Varieties and Constructing Mirror Manifolds*, The Fano Conference, (2004)109-122.
- [Bat00] V.V. Batyrev, *Birational Calabi-Yau n-folds have equal Betti numbers*, New Trends in Algebraic Geometry, (Warwick 2000)1-11.
- [Bat94] V.V. Batyrev, *Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties*, J. Algebraic Geometry., **3** (1994),493-535.
- [BCFKvS00] V. Batrev, I. Ciocan-Fontanie, B. Kim, and D. van Straten, *Mirror symmetry and toric degenerations of partial flag manifolds*, Acta Math. **184** (2000), no. 1, 1-39.
- [BZ01] A. Berenstein and A. Zelevinsky, *Tensor product multiplicities, canonical bases and totally positive varieties*, Invent. Math. **143**(2001),no. 1, 77-128.
- [BZ93] A. Berenstein and A. Zelevinsky, *String bases for quantum groups of type A_r* , Adv. in Soviet Math. **16**, Part 1 (1993),51-89.
- [Cal02] P. Caldero, *Toric degenerations of Schubert varieties*, Transform. Groups **7** (2002), no. 1, 51-60.
- [Ful93] W. Fulton, *Introduction to toric varieties*, Annals of Mathematical Studies **131**, The William H. Roever Lectures in Geometry, Princeton University Press, Princeton, NJ, 1993.
- [GP00] O. Gleizer and A. Postnikov, *Littlewood-Richardson coefficients via Yang-Baxter equations*, Internat. Math. Res. Notices **14** (2000)741-774.
- [GL96] N. Gonciulea and V. Lakshmibai, *Degenerations of flag and Schubert varieties to toric varieties*, Transform. Groups **1** (1996), no. 3, 215-248.
- [K95] M. Kontsevich, Lecture at Orsay (December 7, 1995)
- [Lit98] P. Littelmann, *Cones, crystals, and patterns*, Transform. Groups **3** (1998), no. 2, 145-179.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, ATHENS GA 30602, USA EMAIL: RUSINKO@MATH.UGA.EDU